

Almansi Theorems in Umbral Clifford Analysis and the Quantum Harmonic Oscillator

Guangbin Ren *

University of Science and Technology of China, Department
of Mathematics, Hefei, Anhui 230026, P. R. China

current address: Universidade de Aveiro, Departamento de Matemática

P-3810-193 Aveiro, Portugal

E-mail: ren@ua.pt

Nelson Faustino[†]

Universidade de Aveiro, Departamento de Matemática

P-3810-193 Aveiro, Portugal

E-mail: nfaust@ua.pt

*Research supported by the *Unidade de Investigação “Matemática e Aplicações”* of University of Aveiro, and by the NNSF of China (No. 10771201), NCET-05-0539.

[†]Research supported by *Fundação para a Ciência e a Tecnologia* under the grant SFRH/BD/17657/2004

Almansi Theorems in Umbral Clifford Analysis and the Quantum Harmonic Oscillator

MOS subject classification: 30 G 35, 35 C 10, 39 A 12, 70 H 05

PACS 2006: 02.30.F_n, 03.65.-w

Abstract

We introduce the Umbral calculus into Clifford analysis starting from the abstract of the Heisenberg commutation relation $[\frac{d}{dx}, x] = \mathbf{id}$. The Umbral Clifford analysis provides an effective framework in continuity and discreteness.

In this paper we consider functions defined in a star-like domain $\Omega \subset \mathbb{R}^n$ with values in the Umbral Clifford algebra $C\ell'_{0,n}$ which are Umbral polynomogenic with respect to the (left) Umbral Dirac operator D' , i.e. they belong to the kernel of $(D')^k$. We prove that any polynomogenic function f has a decomposition of the form

$$f = f_1 + x' f_2 + \cdots + (x')^{k-1} f_k,$$

where $x' = x'_1 e_1 + \cdots + x'_n e_n$ and f_j , $j = 1, \dots, k$, are Umbral monogenic functions. As examples, this result recovers the continuous version of the classical Almansi theorem for derivatives and establishes the discrete version of Almansi theorem for difference operator. The approach also provides a similar result in quantum field about Almansi decomposition related to Hamilton operators.

Some concrete examples will be presented for the discrete analog version of Almansi Decomposition and for the quantum harmonic oscillator.

Keywords: Umbral Dirac operator, Almansi's decomposition theorem, Umbral polynomogenic functions, Quantum Harmonic Oscillator.

1 Introduction

We introduce the Umbral Calculus into the Clifford analysis. This provides a framework unifying continuity and discreteness. We verify this phenomena by establishing the Almansi decomposition in Umbral Clifford analysis.

The essence behind continuity and discreteness is Umbral Calculus. Umbral Calculus has many applications in combinatorics, special function theory, approximation theory, statistics, probability, topology and physics (see [1]). Umbral calculus ([2, 3]) can be viewed as an abstract theory of the Heisenberg commutation relation $[P, M] = \mathbf{id}$. In quantum mechanics P is the derivative and M the coordinate operator. The unification of the Umbral structure can be seen by taking the derivative P to be the usual derivative and the divided difference.

In the Umbral Clifford analysis, we take the delta operator O_{x_j} as momentum operators and take $x'_j = \frac{1}{2} \left(x_j O_{x_j}^{-1} + O_{x_j}^{-1} x_j \right)$ as position operators, where $O'_{x_j} f(\underline{x}) = O_{x_j}(x_j f(\underline{x})) - x_j O_{x_j} f(\underline{x})$ is the Pincherle derivative. They satisfy the Heisenberg-Weyl relations

$$[O_{x_j}, O_{x_k}] = 0 = [x'_j, x'_k], \quad [O_{x_j}, x'_k] = \delta_{jk} \mathbf{id}.$$

The Umbral Clifford analysis is a function theory for null solutions for the Umbral Dirac operator

$$D' := \sum_{j=1}^n \mathbf{e}_j O_{x_j}.$$

where $\mathbf{e}_1, \dots, \mathbf{e}_n$ are the Clifford Algebra generators subject to the anti-commuting relations

$$\mathbf{e}_j \mathbf{e}_k + \mathbf{e}_k \mathbf{e}_j = -2\delta_{jk}.$$

By taking O_{x_j} to be the partial derivative $\frac{\partial}{\partial x_j}$, we recover the standard Dirac operator $D = \sum_{j=1}^n \mathbf{e}_j \frac{\partial}{\partial x_j}$. The Clifford analysis is a refinement of harmonic analysis and a generalization of complex analysis to higher dimensions (see [4, 5, 6, 7]). In the spherical harmonics, the Fischer decomposition of polynomials plays a central role. The Almansi decomposition generalized the Fischer decomposition from polynomials to polyharmonic functions. The polyharmonic theory has many applications in elasticity [9], radar imaging [10], and approximation [11, 12].

Almansi's Theorem (cf. [13, 14]) *If f is polyharmonic of degree k in a starlike domain with center 0, then there exist uniquely defined functions $f_1, \dots, f_k$, each harmonic in Ω such that*

$$f(x) = f_1(x) + |x|^2 f_2(x) + \dots + |x|^{2(k-1)} f_k(x).$$

One can find important applications and generalizations of this result in the case of several complex variables in the monograph of ARONSZAJN, CREESE AND LIPKIN, [14], e.g. concerning functions holomorphic in the neighborhood of the origin in $\mathbb{C}^n$. More recent generalizations of Almansi's Theorem can be found in [15, 16, 17, 18].

In [19], Malonek and Ren obtained the Almansi decomposition in $\mathcal{Cl}_{0,n}$. In this paper, we establish the Almansi decomposition in the setting of Umbral Clifford analysis. As a corollary obtained a discrete version of Almansi decomposition. The approach also provides a similar result in quantum field about Almansi decomposition related to Hamilton operators.

2 Umbral calculus revised

2.1 Shift-Invariant Operators

Let $\underline{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$. A monomial over $\underline{x}$ is the product $\underline{x}^\alpha = x_1^{\alpha_1} x_2^{\alpha_2} \dots x_n^{\alpha_n}$. Here and elsewhere $\alpha, \beta, \dots$ denote indices of nonnegative integers. A *polynomial* is a finite linear combination of monomials. We denote the ring of polynomials over $\underline{x}$ by $\mathbb{R}[\underline{x}]$, $\partial_{x_j} := \frac{\partial}{\partial x_j}$ the partial derivative with respect to x_j and $\partial_{\underline{x}} := (\partial_{x_1}, \partial_{x_2}, \dots, \partial_{x_n})$ the gradient operator.

In the sequel, it will be helpful to introduce the shorthand notations

$$\begin{aligned}\partial_{\underline{x}}^\alpha &:= \partial_{x_1}^{\alpha_1} \partial_{x_2}^{\alpha_2} \dots \partial_{x_n}^{\alpha_n}, \\ \alpha! &= \alpha_1! \alpha_2! \dots \alpha_n!, \\ \binom{\beta}{\alpha} &= \frac{\beta!}{\alpha! (\beta - \alpha)!}, \\ |\alpha| &= \sum_{j=1}^n \alpha_j.\end{aligned}$$

Set $\underline{y} \cdot \partial_{\underline{x}} = \sum_{j=1}^n y_j \partial_{x_j}$ and define the shift operator

$$\tau_{\underline{y}} f(\underline{x}) = \exp(\underline{y} \cdot \partial_{\underline{x}}) f(\underline{x}) = f(\underline{x} + \underline{y}).$$

An operator Q is shift-invariant if it commutes with the shift operator $\tau_{\underline{y}}$, when acting on $\mathbb{R}[\underline{x}]$, i.e.

$$[Q, \tau_{\underline{y}}] p(\underline{x}) = 0 \tag{1}$$

holds for all $p \in \mathbb{R}[\underline{x}]$ and $\underline{y} \in \mathbb{R}^n$, where $[\mathbf{a}, \mathbf{b}] = \mathbf{a}\mathbf{b} - \mathbf{b}\mathbf{a}$ denotes the commutator bracket between $\mathbf{a}$ and $\mathbf{b}$. In the language of co-algebra, the shift operator $\tau_{\underline{y}}$ is nothing else than the co-product in a Hopf algebra over $\mathbb{R}[\underline{x}]$, but we shall not discuss this here, see e.g. [20] (Section V.3) for more details.

Furthermore the following multivariate expansion formula holds [21]:

Theorem 2.1 *A linear operator $Q : \mathbb{R}[\underline{x}] \rightarrow \mathbb{R}[\underline{x}]$ is shift-invariant if and only if it can be expressed (as a convergent series) in the gradient $\partial_{\underline{x}}$, that is*

$$Q = \sum_{|\alpha|=0}^{\infty} a_\alpha \partial_{\underline{x}}^\alpha$$

where $a_\alpha = \left[\frac{Q \underline{x}^\alpha}{\alpha!} \right]_{\underline{x}=\underline{0}}$.

We will denote by $\mathcal{U}$ the set of all shift-invariant operators over $\mathbb{R}[\underline{x}]$. This set, endowed with the standard addition and composition between functions, forms an $\mathbb{R}$ -algebra. In the language of umbral calculus, $\mathcal{U}$ is usually named as the Umbral Algebra.

The Pincherle derivative of an operator Q at the coordinate x_j is defined as the commutator

$$Q'_{x_j} = [Q, x_j], \tag{2}$$

where on the right hand side of (2), x_j acts as a multiplication operator on $\mathbb{R}[\underline{x}]$, i.e. $x_j : p(\underline{x}) \mapsto x_j p(\underline{x})$.

According to Theorem 2.1, shift-invariant operators over polynomials are expressed in terms of derivatives. In particular, the multi-index derivatives $\partial_{\underline{x}}^\alpha$ and linear combinations of there are shift-invariant.

Regardless the last viewpoint, one looks to shift-invariant operators as convergent power series $Q(\underline{x}) = \sum_{|\alpha|=0}^{\infty} a_{\alpha} \underline{x}^{\alpha}$, by replacing the vector variable $\underline{x}$ by the gradient $\partial_{\underline{x}}$, namely $\iota[Q(\underline{x})] = Q(\partial_{\underline{x}})$, where $\iota : \mathbb{R}[\underline{x}] \rightarrow \mathcal{U}$ is a mapping between the algebra of polynomials and the umbral algebra. This mapping is one-to-one and onto and therefore an algebra isomorphism.

The inverse image under $\iota : \mathbb{R}[\underline{x}] \rightarrow \mathcal{U}$ of a shift-invariant operator is known as its indicator. As a further consequence, $\partial_{x_j} Q(\underline{x})$ is the indicator of the Pincherle derivative $Q'_{x_j} := Q'_{x_j}(\partial_{\underline{x}})$ c.f. [2]. Therefore, the Pincherle derivative is a derivation operation in the umbral algebra $\mathcal{U}$.

2.2 Quantum Mechanics representation of Umbral Calculus

The main purpose of this subsection is to gather a fully rigorous description of umbral calculus in terms of Boson calculus associated to the second quantization. First we recall some basic facts on Quantum Field Theory.

The Heisenberg-Weyl algebra $\mathcal{F}$ is the free algebra generated from the vacuum vector Φ by the $2n+1$ elements $\mathbf{a}_1^-, \dots, \mathbf{a}_n^-, \mathbf{a}_1^+, \dots, \mathbf{a}_n^+, \mathbf{id}$ satisfying $\mathbf{a}_j \Phi = 0$, $j = 1, \dots, n$ and the commutation relations:

$$[\mathbf{a}_j^-, \mathbf{a}_k^-] = 0 = [\mathbf{a}_j^+, \mathbf{a}_k^+], \quad [\mathbf{a}_j^-, \mathbf{a}_k^+] = \delta_{jk} \mathbf{id}. \quad (3)$$

We now equip $\mathcal{F}$ with an euclidian inner product $\langle \cdot | \cdot \rangle$ such that $\langle \Phi | \Phi \rangle = 1$ and the operators $\mathbf{a}_j^+$ are adjoint to $\mathbf{a}_j^-$:

$$\langle \mathbf{a}_j^+ x | y \rangle = \langle x | \mathbf{a}_j^- y \rangle.$$

The later vector space $(\mathcal{F}, \langle \cdot | \cdot \rangle)$ becomes the Bose algebra, where the elements $\mathbf{a}_j^+$ and $\mathbf{a}_j^-$ (creation and annihilation operators, respectively) forms a basis for the n -particle space.

The explicit expression from the basic vectors in the Fock spaces $\eta_{\alpha} \in \mathcal{F}$ can be derived from the standard lemma of elementary Quantum Field Theory as the multi-index product of creation operators acting on the vacuum vector Φ , i.e.

$$\eta_{\alpha} = \prod_{j=1}^n (\mathbf{a}_j^+)^{\alpha_j} \Phi \quad (4)$$

and moreover, inner products between basic vector fields has the form

$$\langle \eta_{\alpha} | \eta_{\beta} \rangle = \alpha! \delta_{\alpha, \beta}$$

In the following, we will describe the algebra of multivariate polynomials $\mathbb{R}[\underline{x}]$ as a realization of the Bose algebra $\mathcal{F}$.

Among the infinity of possible representations of the simplest one are the multiplication x_j and the derivation operator ∂_{x_j} .

Now consider the multivariate monomials $\underline{x}^{\alpha} \in \mathbb{R}[\underline{x}]$. Evidently x_j and ∂_{x_j} acting on $\underline{x}^{\alpha}$ gives

$$\begin{aligned} x_j \underline{x}^{\alpha} &= \underline{x}^{\alpha + \mathbf{v}_j}, \\ \partial_{x_j} \underline{x}^{\alpha} &= \alpha_j \underline{x}^{\alpha - \mathbf{v}_j} \end{aligned}$$

We now construct the umbral calculus through representations of commutation relations (3) by position and momentum operators, x'_j and O_{x_j} , respectively such that the action of x'_j, O_{x_j} on certain polynomials $V_{\alpha}(\underline{x})$ is analogous to the action of x_j and ∂_{x_j} on the monomials. More specifically we shall search for x'_j, O_{x_j} and their associated polynomials $V_{\alpha}(\underline{x})$ that satisfy

$$x'_j V_{\alpha}(\underline{x}) = V_{\alpha + \mathbf{v}_j}(\underline{x}), \quad O_{x_j} V_{\alpha}(\underline{x}) = \alpha_j V_{\alpha - \mathbf{v}_j}(\underline{x}). \quad (5)$$

These operators can be immediately recognized as raising and lowering operators acting on the polynomials $V_\alpha(\underline{x})$.

In the sequel, we assume that O_{x_j} satisfy the shift-invariance property (1) and $O_{x_j}(x_j)$ is a nonzero constant.

According to the umbral setting such an operator O_{x_j} is called a delta operator and such a polynomial sequence $\{V_\alpha : \alpha \in \mathbb{N}_0^n\}$ generated by (5) is called a *Sheffer set* for the multivariate delta operator $O_{\underline{x}} = (O_{x_1}, O_{x_2}, \dots, O_{x_n})$; the *Sheffer set* $\{V_\alpha : \alpha \in \mathbb{N}_0^n\}$ is called *basic* if $V_\alpha(\mathbf{0}) = 0$ and $V_{\mathbf{0}}(\underline{x}) = \mathbf{1}$. As a further consequence $(O'_{x_j})^{-1}$ always exists and commute with O_{x_j} (c.f. [2], pp. 18).

Using the quantum field representation (4), the *basic* polynomials $V_\alpha(\underline{x})$ are obtained through the action of $(\underline{x}')^\alpha := \prod_{k=1}^n (x'_k)^{\alpha_k}$

$$V_\alpha(\underline{x}) = (\underline{x}')^\alpha \mathbf{1}, \quad (6)$$

which is known as the *Rodrigues formula*.

A further consequence of (5) is the eigenproperty

$$\sum_{j=1}^n x'_j O_{x_j} V_\alpha(\underline{x}) = |\alpha| V_\alpha(\underline{x}) \quad (7)$$

This fact provides a good motivation for calling

$$E' = \sum_{j=1}^n x'_j O_{x_j} \quad (8)$$

the umbral Euler operator, i.e. an operator who measures the degree of V_α ;

The properties of Sheffer-type polynomials are naturally handled within the so called *umbral calculus*. Incidentally we shall study the Sheffer operator

$$\Psi_{\underline{x}} : \underline{x}^\alpha \mapsto V_\alpha(\underline{x}), \quad (9)$$

where $V_\alpha(\underline{x})$ is a Sheffer sequence. This linear operator is obviously invertible and its inverse $\Psi_{\underline{x}}^{-1}$ corresponds to $\Psi_{\underline{x}}^{-1} : V_\alpha(\underline{x}) \mapsto \underline{x}^\alpha$.

This correspondence naturally induces a correspondence between the operators x_j, ∂_{x_j} and O_{x_j}, x'_j . Indeed, according to (5) the linear extension of $\Psi_{\underline{x}} : \underline{x}^\alpha \mapsto V_\alpha(\underline{x})$ is an intertwining operator i.e.

$$O_{x_j} \Psi_{\underline{x}} = \Psi_{\underline{x}} \partial_{x_j}, \quad x'_j \Psi_{\underline{x}} = \Psi_{\underline{x}} x_j. \quad (10)$$

and hence it intertwines the umbral Euler operator (8) and the classical Euler operator $E_{\underline{x}} = \sum_{j=1}^n x_j \partial_{x_j}$, i.e.

$$E' \Psi_{\underline{x}} = \Psi_{\underline{x}} E. \quad (11)$$

Another important consequence of (10) is that $\Psi_{\underline{x}}$ preserves the Heisenberg-Weyl algebra relations (3).

A formal power series is an infinite linear combination of monomials. We will use the variables $\underline{x}$ and $\underline{t}$ to denote formal power series whose coefficients are polynomials and

$$\underline{t} \cdot \underline{x}' = \sum_{j=1}^n t_j x'_j$$

to denote the formal Euclidean inner product between $\underline{t}$ and $\underline{x}'$.

An important consequence of the Heisenberg-Weyl representation (3) concerns the exponential generating function for Sheffer sequences $V_\alpha(\underline{x})$, which is defined by

$$V(\underline{x}, \underline{t}) = \sum_{|\alpha|=0}^{\infty} V_{\alpha}(\underline{x}) \frac{\underline{t}^{\alpha}}{\alpha!} = \sum_{k=0}^{\infty} \frac{1}{k!} (\underline{t} \cdot \underline{x}')^k \mathbf{1} = \exp(\underline{t} \cdot \underline{x}') \mathbf{1}. \quad (12)$$

The above formula corresponds to a formal power series in the variables $\underline{x}$ and $\underline{t}$. According to (10) the exponential generating function (12) may be characterized by means of $V(\underline{x}, \underline{t}) = \Psi_{\underline{x}}(\exp(\underline{t} \cdot \underline{x}))$.

Let us take a close look for the characterization of Sheffer sequences on the umbral algebra.

According to the Theorem 2.1, $\{V_{\alpha} : \alpha \in \mathbb{N}_0^n\}$ is a Sheffer sequence if there is a set of multivariate polynomials $\{O_j : j = 1, \dots, n\}$ such that

$$O_j(\partial_{\underline{x}})V_{\alpha}(\underline{x}) = \alpha_j V_{\alpha - \mathbf{v}_j}(\underline{x}) \quad (13)$$

which are lowering operators. This characterization is not unique, i.e., there are a lot of Sheffer-type sequences satisfying equation (13), for a given $O_j(\underline{x})$. We can further classify them by postulating the existence of the associated raising operator. A general theorem (c.f. [22, 2]) states that a polynomial sequence $\{V_{\alpha} : \alpha \in \mathbb{N}_0^n\}$ satisfying the monomiality principle (5) with a multi-variate operator $O_{\underline{x}} = (O_{x_1}, O_{x_2}, \dots, O_{x_n})$ is uniquely determined by the polynomial vector-field $O(\underline{x}) = (O_1(\underline{x}), O_2(\underline{x}), \dots, O_n(\underline{x}))$ such that

$$O_j(\underline{0}) = \underline{0}, \quad [\partial_{x_j} O_j(\underline{x})]_{\underline{x}=\underline{0}} \neq 0 \quad \text{and} \quad [\partial_{x_j} O_k(\underline{x})]_{\underline{x}=\underline{0}} = 0 \text{ for } j \neq k. \quad (14)$$

The exponential generating function for $V_{\alpha}(\underline{x})$ is then equal to

$$V(\underline{x}, \underline{t}) = \Psi_{\underline{x}}(\exp(\underline{t} \cdot \underline{x})) = \exp(\underline{x} \cdot O^{-1}(\underline{t})) \quad (15)$$

where $O^{-1}(\underline{t}) := (O_1^{-1}(\underline{t}), O_2^{-1}(\underline{t}), \dots, O_n^{-1}(\underline{t}))$.

Furthermore their associated raising and lowering operators associated to (5) are given by

$$O_{x_j} = O_j(\partial_{\underline{x}}), \quad (16)$$

$$x'_j = x_j(O'_{x_j})^{-1}. \quad (17)$$

We would like to emphasize that there many possibilities still for constructing operators x'_j having the raising property. In particular, it is also interesting to consider the special choice

$$x'_j = \frac{1}{2}(x_j(O'_{x_j})^{-1} + (O'_{x_j})^{-1}x_j) \quad (18)$$

Since $(O'_{x_j})^{-1}$ commutes with O_{x_j} (c.f. [2], pp. 18), one can easily verify the Heisenberg-Weyl commutation relations (3) for the operators O_{x_j} and x'_j defined in (16) and (20).

One of major reasons to consider (20) over (17) is that under some special choices of O_{x_j} , the operators O_{x_j} and x'_j becomes symmetric and self-adjoint on a Hilbert space c.f. [29].

Now we would like to point out some properties concerning the umbral operators (16) and (17):

Since O_{x_j} is a delta operator, the operators $(O'_{x_j})^{-1}$ always exists and O'_{x_j} , $(O'_{x_j})^{-1}$ commute with O_{x_j} (c.f. [2], pp. 18), i.e.

$$[O'_{x_j}, O_{x_k}] = 0 = [(O'_{x_j})^{-1}, O_{x_k}] \quad (19)$$

However the same in general does not hold between O'_{x_j} , $(O'_{x_j})^{-1}$ and x'_j defined *viz* formula (17) (see Example 2.1).

We would like also to emphasize that there many possibilities still for constructing operators x'_j having the raising property. In particular, it is also interesting to consider the special choice

$$x'_j = \frac{1}{2}(x_j(O'_{x_j})^{-1} + (O'_{x_j})^{-1}x_j) \quad (20)$$

One can easily verify the Heisenberg-Weyl commutation relations (3) for the operators O_{x_j} and x'_j defined in (16) and (20). On the other hand we also get the raising property for the operators (20). However like for (17), the operators (20) does not commute with O'_{x_j} and $(O'_{x_j})^{-1}$.

In brief, one of major reasons to consider (20) instead of (17) is that under some special choices of O_{x_j} , the operators O_{x_j} and x'_j becomes symmetric and self-adjoint on a Hilbert space c.f. [29].

Here there are some examples of so obtained representations of the Heisenberg-Weyl algebra:

Example 2.1 (Forward/Backward differences) Denote by $\partial_h^{\pm j}$ the forward/backward difference operators by

$$\partial_h^{\pm j} = \pm \frac{\tau_{\pm h \mathbf{v}_j} - \mathbf{id}}{h} = \pm \frac{1}{h} (\exp(\pm h \partial_{x_j}) - \mathbf{id}).$$

These operators mutually commute when acting on functions, i.e.

$$\partial_h^{\pm j} (\partial_h^{\pm k} f(\underline{x})) = \partial_h^{\pm k} (\partial_h^{\pm j} f(\underline{x})). \quad (21)$$

and are interrelated with shifts $\tau_{\pm h \mathbf{v}_j}$

$$\tau_{-h \mathbf{v}_j} (\partial_h^{+j} f)(\underline{x}) = (\partial_h^{-j} f)(\underline{x}), \quad \tau_{h \mathbf{v}_j} (\partial_h^{-j} f)(\underline{x}) = (\partial_h^{+j} f)(\underline{x}).$$

Let us remark that the finite difference action $\partial_h^{\pm j}$ acting on functions satisfies the product rule

$$\partial_h^{\pm j} (x_j f(\underline{x})) = \pm \frac{(x_j \pm h) f(\underline{x} + h \mathbf{v}_j) - x_j f(\underline{x})}{h} = x_j \partial_h^{\pm j} (x_j f(\underline{x})) + f(\underline{x} + h \mathbf{v}_j). \quad (22)$$

and hence $(\partial_h^{\pm j})' = \tau_{\pm h \mathbf{v}_j}$. Replacing the coordinate function x_j by the operators $x_j \tau_{\mp h \mathbf{v}_j}$, we establish the duality between the finite difference operators $\partial_h^{\pm j}$ and the “formal” coordinate functions $x_j \tau_{\pm h \mathbf{v}_j}$.

We also note that the operators $x_j \tau_{\pm h \mathbf{v}_j}$ mutually commute, when acting on functions

$$x_j \tau_{\pm h \mathbf{v}_j} (x_k \tau_{\pm h \mathbf{v}_k} f(\underline{x})) = x_k \tau_{\pm h \mathbf{v}_k} (x_j \tau_{\pm h \mathbf{v}_j} f(\underline{x})). \quad (23)$$

Furthermore, the commutative relations (21), (23) together with the duality relation (22) endow an algebraic representation of the Heisenberg-Weyl (3) algebra, where the “formal” coordinate functions $x_j \tau_{\mp h \mathbf{v}_j}$ represent “creation” operators dual to the “annihilation” operators $\partial_h^{\pm j}$.

Applying the Rodrigues formula (6), we obtain the following multivariate polynomials

$$(x)_{\pm}^{(\alpha)} = \prod_{j=1}^n x_j (x_j \pm h) \dots (x_j \pm (\alpha_j - 1)h).$$

These polynomials are the basic polynomials that satisfy the equations (5). It is interesting to see that these multivariate polynomials coincide with the multi-index factorial powers $(x)_{\pm}^{(\alpha)}$ introduced by N. Faustino and U. Kähler in [25].

Here we would like to emphasize that the operators $x_j \tau_{\pm h \mathbf{v}_j}$ and $\tau_{\pm h \mathbf{v}_j}$ does not commute when acting on functions, i.e.

$$\begin{aligned} \tau_{\pm h \mathbf{v}_j} (x_k \tau_{\pm h \mathbf{v}_k} f(\underline{x})) &= \pm h \delta_{jk} f(\underline{x} \pm h(\mathbf{v}_j + \mathbf{v}_k)) + x_k \tau_{\pm h \mathbf{v}_k} (\tau_{\pm h \mathbf{v}_j} f(\underline{x})), \\ \tau_{\mp h \mathbf{v}_j} (x_k \tau_{\pm h \mathbf{v}_k} f(\underline{x})) &= \mp h \delta_{jk} f(\underline{x} \pm h(\mathbf{v}_k - \mathbf{v}_j)) + x_k \tau_{\pm h \mathbf{v}_k} (\tau_{\pm h \mathbf{v}_j} f(\underline{x})) \end{aligned}$$

Another family of polynomials involving the position operator x'_j defined in (20) can be considered. In particular, recursive application of the Rodrigues formula (6) yields the following family of polynomials.

$$\prod_{j=1}^n \frac{1}{2^{\alpha_j}} (x_j \tau_{\pm h \mathbf{v}_j} + \tau_{\pm h \mathbf{v}_j} x_j)^{\alpha_j} \mathbf{1} = \prod_{j=1}^n \left(x_j \pm \frac{1}{2}h \right) \left(x_j \pm \frac{3}{2}h \right) \dots \left(x_j \pm \frac{2\alpha_j - 1}{2}h \right).$$

We will come back to the Heisenberg-Weyl character of the operators $\partial_h^{\pm j}$ and x'_j in Section 6 when establishing the comparison between the Umbral version of Almans-type decomposition and the Fischer Decomposition for Difference Dirac operators obtained in [25].

Example 2.2 (Central difference operator) In lattice gauge theories it is common to consider the central finite difference operator (i.e. the average between the forward and backward differences)

$$O_{x_j} = \frac{1}{2}(\partial_h^{+j} + \partial_h^{-j}) = \frac{1}{h} \sinh(h\partial_{x_j})$$

instead of the forward/backward differences $\partial_h^{\pm j}$.

One among many reasons behind this choice is that contrary to $\partial_h^{\pm j}$, O_{x_j} are symmetric and self-adjoint on the Hilbert space $\ell_2(h\mathbb{Z}^n)$.

Furthermore the Pincherle derivative for O_{x_j} corresponds to

$$O'_{x_j} = \frac{\tau_{h\mathbf{v}_j} + \tau_{-h\mathbf{v}_j}}{2} = \cosh(h\partial_{x_j}).$$

It can be easily seen that the multivariate operator $O_{\underline{x}} = (O_{x_1}, O_{x_2}, \dots, O_{x_n})$ is a delta operator so it admits a basic polynomial sequence satisfying the equations (5) and that O'_{x_j} has an inverse since $O'_{x_j} 1 = 1 \neq 0$.

Now it arises the question how to compute $(O'_{x_j})^{-1}$. Since the map $\iota : \mathbb{R}[\underline{x}] \rightarrow \mathcal{U}$ is an algebra isomorphism, the problem of finding an inverse for O'_{x_j} it is enough to find an inverse for the series expansion

$$\cosh(hx_j) = \sum_{k=0}^{\infty} \frac{1}{(2k)!} (hx_j)^{2k}$$

We however note that finding an inverse for $\cosh(hx_j)$ becomes increasingly cumbersome. Nevertheless, there is a certain systematic in it.

For a sake of simplicity, we impose periodic boundary conditions on our lattice such that $\underline{x} + N\mathbf{v}_j = \underline{x}$, i.e. $(\tau_{h\mathbf{v}_j})^N = \text{id}$ for a certain $N \in \mathbb{N}$. (i.e. a periodic lattice). In particular, when $N/2$ is odd, $(O'_{x_j})^{-1}$ corresponds to c.f. [29]

$$(O'_{x_j})^{-1} = \sum_{k=0}^{N/2-1} (-1)^k (\tau_{h\mathbf{v}_j})^k$$

while for N odd one obtain

$$(O'_{x_j})^{-1} = \sum_{k=0}^{(N-1)/2} (-1)^{k+(N-1)/2} (\tau_{h\mathbf{v}_j})^{2k} + \sum_{k=0}^{(N-1)/2-1} (-1)^k (\tau_{h\mathbf{v}_j})^{2k+1}.$$

These inversion formulae will be of special interest on Section 6 when we consider the gauged version of the Harmonic oscillator.

3 Almansi-type theorems in Umbral Clifford analysis

3.1 Umbral Clifford Analysis

The main objective of this subsection is to introduce a well-adapted Clifford setting.

In what follows we retain the notation adopted in subsection 2.2. Additionally we will denote by $\mathcal{Cl}_{0,n}$ the algebra of signature $(0, n)$ whose generators $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$ satisfy the graded deformed anti-commuting relations

$$\{\mathbf{e}_j, \mathbf{e}_k\} = -2\delta_{jk}, \quad (24)$$

where $\{\mathbf{a}, \mathbf{b}\} = \mathbf{ab} + \mathbf{ba}$ denotes the bracket between $\mathbf{a}$ and $\mathbf{b}$.

The following algebra is an algebra of radial type, i.e.

$$\{ [\mathbf{e}_j, \mathbf{e}_k], \mathbf{e}_l \} = 0, \quad \text{for all } j, k, l = 1, \dots, n. \quad (25)$$

In the above definition there is indeed no a priori defined a linear space to which the generators $\mathbf{e}_j$ belong. Nevertheless, by only using (25) one can already deduce many properties [23].

Analogously to [23], we define Umbral Clifford analysis as the study of the operators belonging to the algebra of differential operators

$$\text{Alg} \{ x'_j, O_{x_j}, \mathbf{e}_j : j = 1, \dots, n \}, \quad (26)$$

where the operators O_{x_j} and x'_j are defined accordingly to (16) and (17)/(20), respectively.

We define the umbral counterpart for the Dirac operator as

$$D' = \sum_{j=1}^n \mathbf{e}_j O_{x_j}. \quad (27)$$

and by

$$x' = \sum_{j=1}^n \mathbf{e}_j x'_j \quad (28)$$

the umbral counterpart of the vector variable.

The Heisenberg-Weyl character of the operators x'_j and O_{x_j} together with the commutation relations (19), and the anti-commutation rule (24) implies that

$$(x')^2 = - \sum_{j=1}^n (x'_j)^2, \quad (29)$$

$$(D')^2 = - \sum_{j=1}^n O_{x_j}^2. \quad (30)$$

The operator $(x')^2$ then corresponds to the generalization of the norm squared of a vector in the Euclidean space since $|x'|^2 = -(x')^2 \mathbf{1}$ while the right hand side of (30) corresponds to the umbral counterpart of the Laplacian, i.e. $\Delta' = -(D')^2$, and this indeed what we expected from a "true" coordinate variable and Dirac operator.

Bellow there are some examples of this type of operators:

Example 3.1 If $O_{x_j} = \partial_{x_j}$, we recover the standard Clifford Analysis since the umbral operators D' and x' coincide with the standard Dirac and coordinate variable operators, respectively. Furthermore the continuum Hamiltonian $\frac{1}{2}(-\Delta + |\underline{x}|^2)$ can be rewritten as

$$\frac{1}{2}(-\Delta + |\underline{x}|^2) = \frac{1}{2}(D^2 - x^2),$$

where the quantity $\frac{1}{2}|\underline{x}|^2 = -\frac{1}{2}x^2$ should be interpreted as a spherical symmetric potential operator.

Example 3.2 If $O_{x_j} = \partial_h^{+j}$, where ∂_h^{+j} are the forward differences introduced on Example 2.1, the corresponding operator D' is given by

$$D' = \sum_{j=1}^n \mathbf{e}_j \partial_h^{\pm j}. \quad (31)$$

The square of $(D')^2$ does not split the star laplacian

$$\Delta' \neq \sum_{j=1}^n \partial_h^{+j} \partial_h^{-j}, \quad (32)$$

Example 3.3 If $O_{x_j} = \frac{1}{2} (\partial_h^{+j} + \partial_h^{-j})$ are the central differences introduced on Example 2.2, the corresponding operator D' is given by

$$D' = \frac{1}{2} \sum_{j=1}^n \mathbf{e}_j (\partial_h^{+j} + \partial_h^{-j}). \quad (33)$$

The square of $(D')^2$ split the star laplacian

$$\Delta' = \sum_{j=1}^n \partial_{2h}^{+j} \partial_{2h}^{-j}, \quad (34)$$

In Examples 3.2 and 3.3, the operator $-\frac{1}{2}(x')^2$ describes a spherical potencial of the lattice while the equation $-(x')^2 \mathbf{1} = r$ describes a lattice sphere on the n -dimensional ambient space.

We would like to point out that under the conditions of (2.2), the special choice (20) leads to spherical potentials on the lattice which possesses the symmetric property like in *continuum*. However the same does not holds under the conditions of Example 2.1.

We will come back to this problem in Section 6 when we consider the gauged version of the harmonic oscillator.

3.2 Decomposition of Umbral polynomogonic functions

Definition 3.1 Let Ω be a domain in $\mathbb{R}^n$ and $k \in \mathbb{N}$. A function $f : \Omega \longrightarrow \mathcal{Cl}_{0,n}$ is Umbral polynomogonic of degree k if $(D')^k f = 0$. If $k = 1$, it is called Umbral monogenic function.

Definition 3.2 A domain $\Omega \subset \mathbf{R}^n$ is a starlike domain with center 0 if with $x \in \Omega$ also $tx \in \Omega$ holds for any $0 \leq t \leq 1$.

Let Ω be a starlike domain with center 0. For any $s > 0$, we define the operator $I_s : C(\Omega, \mathcal{Cl}_{0,n}) \longrightarrow C(\Omega, \mathcal{Cl}_{0,n})$ by

$$I_s f(x) = \int_0^1 f(tx) t^{s-1} dt. \quad (35)$$

Recalling the definition of $\Psi_{\underline{x}}$ in (9), we now introduce an operator

$$I'_s = \Psi_{\underline{x}} I_s \Psi_{\underline{x}}^{-1}. \quad (36)$$

For any $k \in \mathbb{N}$, denote

$$Q'_k = \frac{1}{c_k} I'_{\frac{n}{2}} I'_{\frac{n}{2}+1} \cdots I'_{\frac{n}{2} + [\frac{k-1}{2}]}, \quad (37)$$

where $c_k = (-2)^k [k/2]!$.

Theorem 3.1 *Let Ω be a starlike domain in $\mathbb{R}^n$ with center 0. If f is a Umbral polymonogenic function in Ω of degree k , then there exist unique functions $f_1, \dots, f_k$, each Umbral monogenic in Ω , such that*

$$f(x) = f_1(x) + x'f_2(x) + \dots + (x')^{k-1}f_k(x). \quad (38)$$

Moreover the Umbral monogenic functions $f_1, \dots, f_k$ are given by the the following formulas:

$$\begin{aligned} f_k &= Q'_{k-1}(D')^{k-1}f(x) \\ f_{k-1} &= Q'_{k-2}(D')^{k-2}(\mathbf{id} - (x')^{k-1}Q'_{k-1}(D')^{k-1})f(x) \\ &\vdots \\ f_2 &= Q_1D'(\mathbf{id} - (x')^2Q_2(D')^2) \dots (\mathbf{id} - (x')^{k-1}Q_{k-1}(D')^{k-1})f(x) \\ f_1 &= (\mathbf{id} - x'Q'_1D')(\mathbf{id} - (x')^2Q'_2(D')^2) \dots (\mathbf{id} - (x')^{k-1}Q_{k-1}(D')^{k-1})f(x). \end{aligned} \quad (39)$$

Conversely, the sum in (38), with $f_1, \dots, f_k$ Umbral monogenic in Ω , defines a Umbral polymonogenic function in Ω .

3.3 The proof of the Almansi Theorem

Before proving Theorem 3.1, we need some lemmas.

Denote

$$E_s = s\mathbf{id} + E. \quad (40)$$

We write E instead of E_0 when $s = 0$.

Lemma 3.1 ([19]) *Let $x \in \mathbb{R}^n$ and Ω be a domain with $\Omega \supset [0, x]$. If $s > 0$ and $f \in C^1(\Omega, \mathcal{Cl}'_{0,n})$, then*

$$f(x) = I_s E_s f(x) = E_s I_s f(x). \quad (41)$$

Note that Lemma 3.1 remains true in the setting of Umbral Clifford analysis, since it holds componentwise.

Since $I'_s = \Psi_{\underline{x}} I_s \Psi_{\underline{x}}^{-1}$ and $E'_s = \Psi_{\underline{x}} E_s \Psi_{\underline{x}}^{-1}$, it follows from (41) that

$$E'_s I'_s = I'_s E'_s = \mathbf{id}.$$

Lemma 3.2 *Let Ω be a domain in $\mathbb{R}^n$. If $f \in C^1(\Omega)$, then*

$$D'E'_s f(x) = E'_{s+1} D' f(x), \quad D' I'_s f(x) = I'_{s+1} D' f(x). \quad (42)$$

Proof: Recall that $[O_{x_j}, x'_k] = \delta_{jk}\mathbf{id}$ and $[O_{x_j}, O_{x_k}] = 0$, i.e.,

$$O_{x_j} x'_k = \delta_{jk}\mathbf{id} + x'_k O_{x_j}, \quad O_{x_j} O_{x_k} = O_{x_k} O_{x_j}.$$

By definition

$$D' = \sum_{j=1}^n e_j O_{x_j}, \quad E' = \sum_{k=1}^n x'_k O_{x_k},$$

so that

$$\begin{aligned} D'E' &= \sum_{j,k=1}^n e_j O_{x_j} x'_k O_{x_k} \\ &= \sum_{j,k=1}^n e_j (\delta_{jk}\mathbf{id} + x'_k O_{x_j}) O_{x_k} \\ &= \sum_{j,k=1}^n \delta_{jk} e_j O_{x_k} + \sum_{j,k=1}^n e_j x'_k O_{x_j} O_{x_k} \\ &= \sum_j e_j O_{x_j} + \sum_{j,k=1}^n x'_k O_{x_k} e_j O_{x_j} \\ &= D' + E'D' \\ &= E'_1 D'. \end{aligned}$$

With the same approach, we can also show that $x'D' + D'x' = -2E'_{n/2}$ and $E'_s x' = x' E'_{s+1}$.

Hence $D'E'_s = D'(s\mathbf{id} + E') = sD' + E'_1 D' = E'_{s+1} D'$. Since $E'_s = I'^{-1}_s$, we have

$$D'I'_s = I'_{s+1} E'_{s+1} D'I'_s = I'_{s+1} D'E'_s I'_s = I'_{s+1} D'.$$

■

As direct consequence of (42), we find that $I'_s f$ and $E'_s f$ are monogenic, whenever f is Umbral monogenic. From the definition of Q'_k in (37), we thus obtain

$$Q'_k H_1 = H_1. \quad (43)$$

Lemma 3.3 *Let Ω be a starlike domain in $\mathbb{R}^n$. For any monogenic function f in Ω ,*

$$(D')^k (x')^k Q'_k f(x) = f(x), \quad x \in \Omega.$$

Proof: From (37) and Lemma 3.2, we know that

$$(Q'_k)^{-1} = c_k E'_{\frac{n}{2} + [\frac{k-1}{2}]} \cdots E'_{\frac{n}{2}+1} E'_{\frac{n}{2}}. \quad (44)$$

Denote $g = Q'_k f$. From (43) and (44), g is Umbral monogenic in Ω and

$$f(x) = c_k E'_{\frac{n}{2} + [\frac{k-1}{2}]} \cdots E'_{\frac{n}{2}+1} E'_{\frac{n}{2}} g(x).$$

We need to show

$$(D')^k (x')^k g(x) = c_k E'_{\frac{n}{2} + [\frac{k-1}{2}]} \cdots E'_{\frac{n}{2}+1} E'_{\frac{n}{2}} g(x), \quad (45)$$

or rather

$$\begin{aligned} (D')^{2m-1} (x'^{2m-1}) g(x) &= -(m-1)! 2^{2m-1} E'_{\frac{n}{2}+m-1} \cdots E'_{\frac{n}{2}+1} E'_{\frac{n}{2}} g(x), \\ (D')^{2m} (x'^{2m}) g(x) &= m! 2^{2m} E'_{\frac{n}{2}+m-1} \cdots E'_{\frac{n}{2}+1} E'_{\frac{n}{2}} g(x) \end{aligned} \quad (46)$$

for any g Umbral monogenic in Ω and $m \in \mathbb{N}$.

We use induction to prove (46). Since $x'D' + D'x' = -2E'_{n/2}$ and $Dg(x) = 0$, we have

$$D'(x'g(x)) = -2E'_{\frac{n}{2}} g(x). \quad (47)$$

Next we show that, for any $x \in \Omega$ and $m \in \mathbb{N}$,

$$\begin{aligned} D'((x')^{2m} g(x)) &= -2m(x')^{2m-1} g(x); \\ D'((x')^{2m-1} g(x)) &= -2(x')^{2(m-1)} E'_{\frac{n}{2}+m-1} g(x). \end{aligned} \quad (48)$$

This can be checked by induction. Assuming that (48) holds for m , we shall now prove it also holds for $m+1$. We now apply the operator $x'D' + D'x' = -2E'_{n/2}$ to the function $(x')^{2m} g(x)$:

$$x'D'((x')^{2m} g(x)) + D'x'((x')^{2m} g(x)) = -2E'_{n/2}((x')^{2m} g(x)). \quad (49)$$

By the hypothesis of induction, the first term in the left is equal to $-2m(x')^{2m} g(x)$, while the left side equals $-2x'^{2m} E'_{n/2+2m} g(x)$ due to the fact that

$$E'_s x' = x' E'_{s+1}.$$

As a result,

$$\begin{aligned} D'(x'^{2m+1} g(x)) &= -x'D'(x'^{2m} g(x)) - 2E'_{n/2}(x'^{2m} g(x)) \\ &= 2mx'^{2m} g(x) - 2x'^{2m} E'_{n/2+2m} g(x) \\ &= -2x'^{2m} E'_{n/2+m} g(x). \end{aligned} \quad (50)$$

This proves the second equality of (48). The first equality of (48) can be proved similarly. This proves the identities (48).

Finally we apply (48) to prove (46). It is easy to prove when $m = 1$. For the general case, from (48) we have

$$\begin{aligned} (D')^{2m}((x')^{2m}g(x)) &= (D')^{2m-1}D'((x')^{2m}g(x)) \\ &= -2m(D')^{2(m-1)}((x')^{2m-1}g(x)) \end{aligned}$$

and

$$\begin{aligned} (D')^{2m-1}((x')^{2m-1}g(x)) &= (D')^{2(m-1)-1}D'(D'((x')^{2m-1}g(x))) \\ &= (D')^{2(m-1)-1}D'(-2(x')^{2(m-1)}E'_{\frac{n}{2}+m-1}g(x)) \\ &= 4(m-1)(D')^{2(m-1)-1}((x')^{2(m-1)-1}E'_{\frac{n}{2}+m-1}g(x)). \end{aligned}$$

(46) follows directly from the above induced formulas and the assumption of induction. ■

Now we come to the proof of our main theorem.

Proof of Theorem 3.1: Recall $H_k = \{f \in C^\infty(\Omega, \mathcal{C}l_{0,n}) : (D')^k f = 0\}$. It is sufficient to show that

$$H_k = H_{k-1} + T_{k-1}H_1, \quad k \in \mathbb{N},$$

where $T_k = (x')^k$. Notice that Lemma 3.3 states that

$$(D')^k T_k Q'_k = \text{id}. \quad (51)$$

We divide the proof into two parts.

(i) $H_k \supset H_{k-1} + T_{k-1}H_1$.

Since $H_{k-1} \subset H_k$, we need only to show $T'_{k-1}H_1 \subset H_k$. For any $g \in H_1$, by (51) and (43) we have

$$(D')^k (T_{k-1}g) = D'((D')^{k-1}T_{k-1}Q_{k-1})(Q'_{k-1})^{-1}g = D'(Q'_{k-1})^{-1}g = 0.$$

(ii) $H_k \subset H_{k-1} + T_{k-1}H_1$.

For any $f \in H_k$, we have the decomposition

$$f = (\text{id} - T'_{k-1}Q'_{k-1}D'^{k-1})f + T_{k-1}(Q'_{k-1}(D')^{k-1}f).$$

We will show that the first summand above is in H_{k-1} and the item in the braces of the second summand is in H_1 . This can be verified directly. First,

$$\begin{aligned} (D')^{k-1}(\text{id} - T_{k-1}Q_{k-1}(D')^{k-1})f &= ((D')^{k-1} - ((D')^{k-1}T_{k-1}Q'_{k-1})(D')^{k-1})f \\ &= ((D')^{k-1} - (D')^{k-1})f = 0. \end{aligned}$$

Next, since $(D')^{k-1}f \in H_1$ and $Q'_{k-1}H_1 \subset H_1$, we have $Q'_{k-1}(D')^{k-1}f \in H_1$, as desired.

This proves that $H_k = H_{k-1} + T_{k-1}H_1$. By induction, we can easily deduce that $H_k = H_1 + T_1H_1 + \dots + T_{k-1}H_1$.

Next we prove that for any $f \in H_k$ the decomposition

$$f = g + T_{k-1}f_k, \quad g \in H_{k-1}, f_k \in H_1$$

is unique. In fact, for such a decomposition, applying $(D')^{k-1}$ on both sides we obtain

$$\begin{aligned} (D')^{k-1}f &= (D')^{k-1}g + (D')^{k-1}T_{k-1}f_k \\ &= (D')^{k-1}T_{k-1}Q'_{k-1}(Q'_{k-1})^{-1}f_1 \\ &= (Q'_{k-1})^{-1}f_k. \end{aligned}$$

Therefore

$$f_k = Q'_{k-1}(D')^{k-1}f,$$

so that

$$g = f - T_{k-1}f_k = (\mathbf{id} - T_{k-1}Q'_{k-1}(D')^{k-1})f.$$

Thus (68) follows by induction.

To prove the converse, we see from (45) that, for any $k \in \mathbb{N}$, $(D')^{k+1}(x')^k H_1 = 0$. Replacing k by j , we have

$$(D')^k(x')^j H_1 = 0$$

for any $k > j$. ■

4 The Quantum Harmonic Oscillator

In this section, for a set of creation and annihilation operators $\mathbf{a}_j^+$ and $\mathbf{a}_j^-$, respectively, we shall decompose the Hamiltonian of the classical harmonic oscillator

$$\mathcal{H} = \frac{1}{2} \sum_{j=1}^n -(\mathbf{a}_j^-)^2 + (\mathbf{a}_j^+)^2$$

into two first order operators D_+ and D_- . The operators D_+ and D_- constitute a pair of the same roles as a vector operator involving position and a momentum operators, respectively. We thus can establish the Almansi decomposition with respect to this pair similar to our main result in the Almansi decomposition for the ordinary pair.

4.1 Classical Hamiltonian splitting and Umbral Calculus

We first consider the classical case. We set the Hamiltonian operator $\mathcal{H} = \frac{1}{2}(-\Delta + |\underline{x}|^2)$ and put $\mathbf{a}_j^\pm = \frac{1}{\sqrt{2}}(x_j \mp \partial_{x_j})$.

As the operators $\mathbf{a}_j^\pm$ satisfy the Heisenberg-Weyl commutation relations (3), we can split $\mathcal{H}$ as

$$\mathcal{H} = E^{+-} + \frac{n}{2}\mathbf{id},$$

where $E^{+-} := \sum_{j=1}^n \mathbf{a}_j^+ \mathbf{a}_j^-$ is Hamiltonian of a field of free (non-interacting) bosons. This operator plays the same role of the Umbral Euler operator (8).

For $D = \sum_{j=1}^n \mathbf{e}_j \partial_{x_j}$ and $x = \sum_{j=1}^n \mathbf{e}_j x_j$, where $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$ are the standard Clifford generators satisfying the anti-commuting relations

$$\{\mathbf{e}_j, \mathbf{e}_k\} = -2\delta_{jk}\mathbf{id},$$

we set $D_\pm = \frac{1}{\sqrt{2}}(x \mp D) = \sum_{j=1}^n \mathbf{e}_j \mathbf{a}_j^\mp$. Hence the operators $D_\pm$ and $\mathcal{H}$ satisfy the following relations

$$\{D_+, D_-\} = -2\mathcal{H}, \quad [H, D_-] = -D_-, \quad [H, D_+] = D_+. \quad (52)$$

Using the above relations one may easily calculate the following commutators

$$[D_+^2, D_-^2] = 4\mathcal{H}, \quad [\mathcal{H}, D_-^2] = -2D_-^2, \quad [\mathcal{H}, D_+^2] = 2D_+^2, \quad (53)$$

$$[D_+^2, D_+] = 0, \quad [D_-^2, D_+] = 2D_-, \quad [\mathcal{H}, D_+^2] = 2D_+, \quad (54)$$

$$[D_-, D_+^2] = -2D_+. \quad (55)$$

From the relations (53), we conclude that $\frac{1}{2}D_+^2$, $\frac{1}{2}D_-^2$ and $\mathcal{H}$ are the canonical generators of the Lie algebra $sl_2(\mathbb{R})$ and hence we have a representation of harmonic analysis for the classical Harmonic oscillator (see e.g. [8]). On the other hand, there is such a kind of Clifford Analysis framework involving the operators D_+ and D_- since

$$\text{span} \left\{ \frac{1}{2}D_-^2, \frac{1}{2}D_+^2, \mathcal{H} \right\} \oplus \text{span} \{D_-, D_+\}$$

equipped with the standard graded commutator $[\cdot, \cdot]$ is isomorphic to a Lie superalgebra of type $osp(1|2)$ (see e.g. [27]).

5 Almansi-type decomposition for the Umbral Harmonic Oscillator

Let us consider the umbral counterpart of the Hamiltonian written in the language of Umbral Clifford Analysis corresponds to

$$\mathcal{H}' = \frac{1}{2} ((D')^2 - (x')^2).$$

where D' and x' are Umbral Dirac and the Umbral coordinate variable defined in equations (27) and (28), respectively. Next we put

$$D'_\pm := \frac{1}{\sqrt{2}}(x' \mp D') \quad (56)$$

The representations of the Lie algebra $sl_2(\mathbb{R})$ and the Lie superalgebra $osp(1|2)$ also fulfil in the umbral setting. Indeed, if we consider the Sheffer operator $\Psi_{\underline{x}}$ defined in (9), the umbral operators

$$\frac{1}{\sqrt{2}}(x'_j \mp O_{x_j}) = \Psi_{\underline{x}} \mathbf{a}_j^\pm \Psi_{\underline{x}}^{-1}$$

satisfy the Heisenberg-Weyl commutation relations (3).

Moreover since $D'_\pm = \Psi_{\underline{x}} D_\pm \Psi_{\underline{x}}^{-1}$ and $\mathcal{H}' = \Psi_{\underline{x}} \mathcal{H} \Psi_{\underline{x}}^{-1}$, the relations (52)-(55) also fulfils for the operators $D'_\pm$, $(D'_\pm)^2$ and $\mathcal{H}'$. Now we are in conditions to formulate the Almansi-type decomposition for the Umbral Harmonic Oscillator.

Here the main step consists in replace the Umbral Dirac operator D' by D'_- and the umbral coordinate vector variable x' by D'_+ .

We take the definition of Ψ_x , I'_s , Q'_k as in (9), (36) and (37) by replacing the operators x'_j , E'_s and I'_s by $\mathbf{a}_j^+$, $(s - \frac{n}{2})\mathbf{id} + \mathcal{H}'$ and $((s - \frac{n}{2})\mathbf{id} + \mathcal{H}')^{-1}$, respectively.

Hence we can deduce that if $D'_-g(x) = 0$ then

$$D'_-((D'_+)^{2m}g(x)) = -2m(D'_+)^{2m-1}g(x), \quad D'_-((D'_+)^{2m-1}g(x)) = -2(D'_+)^{2(m-1)}g(x);$$

Moreover

$$D'^k_-((D'_+)^k g(x)) = g(x).$$

Let Ω be a domain in $\mathbb{R}^n$ and $k \in \mathbb{N}$ and let $H_k^-(\Omega)$ denote the set of all functions $f : \Omega \longrightarrow \mathcal{Cl}_{0,n}$ such that $(D'_-)^k f = 0$. With the same proof as in Theorem 3.1, we have the following Almansi decomposition in terms the eigenvectors of the Umbral Hamiltonian $\mathcal{H}'$.

Theorem 5.1 *Let Ω be a starlike domain in $\mathbb{R}^n$ with center 0. If $f \in H^-(\Omega)$, then there exist unique functions $f_1, \dots, f_k$, each in $H_1^-(\Omega)$, such that*

$$f(x) = f_1(x) + D'_+ f_2(x) + \dots + (D'_+)^{k-1} f_k(x). \quad (57)$$

Moreover $f_1, \dots, f_k$ are given by the the following formulas:

$$\begin{aligned} f_k &= Q'_{k-1}(D'_-)^{k-1} f(x) \\ f_{k-1} &= Q'_{k-2}(D'_-)^{k-2} (\mathbf{id} - (D'_+)^{k-1} Q'_{k-1}(D'_-)^{k-1}) f(x) \\ &\vdots \\ f_2 &= Q'_1 D'_- (\mathbf{id} - (D'_+)^2 Q'_2 (D'_-)^2) \dots (\mathbf{id} - (D'_+)^{k-1} Q'_{k-1}(D'_-)^{k-1}) f(x) \\ f_1 &= (\mathbf{id} - D'_+ Q'_1 D'_-)(\mathbf{id} - (D'_+)^2 Q'_2 (D'_-)^2) \dots (\mathbf{id} - (D'_+)^{k-1} Q'_{k-1}(D'_-)^{k-1}) f(x). \end{aligned} \quad (58)$$

Conversely, the sum in (57), with $f_1, \dots, f_k \in H_1^-(\Omega)$, defines a function in $H_k^-(\Omega)$.

6 Umbral Bridge

We can now show that the powerful "Umbral bridge" connects continuity and discreteness.

First we consider the continuous case. Namely, we take O_{x_j} to be the usual continuous derivative ∂_{x_j} . Then the Pincherle derivative $(\partial_{x_j})' = [\partial_{x_j}, x_j] = \mathbf{id}$, D' is exactly the classical Dirac operator, so that the resulting Umbral Clifford algebra turns out to be the usual Clifford algebra $\mathcal{Cl}_{0,n}$ and hence our main theorem then recoveries the Almansi decomposition as in [19].

It is now natural to take a look for applications of the umbral calculus formalism in problems related to lattices. Our main objective in this section is therefore to find a Almansi-type decomposition for the gauged version of the Dirac Operator and for the quantum harmonic oscillator.

First we will explain how to recover the Almansi-type decomposition in the discrete case.

Example 6.1 (Discrete counterpart of Almansi Decomposition) *Our starting point is to consider the difference Dirac operators $D_h^\pm$ introduced in [25]. In the terminology of that paper,*

$$D_h^\pm = \sum_{j=1}^n \mathbf{e}_j \partial_h^{\pm j}$$

are the forward/backward versions of the Dirac operator, where $\partial_h^{\pm j}$ introduced in Example 2.1 and $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$ being the standard basis for the Clifford Algebra $\mathcal{Cl}_{0,n}$.

By introducing the Sheffer maps

$$\Psi_\pm^\pm : \underline{x}^\alpha \mapsto (\underline{x})_\mp^{(\alpha)},$$

the Almansi-type decomposition for the operators can be obtained by replacing the standard Dirac $D = \sum_{j=1}^n \mathbf{e}_j \partial_{x_j}$ and coordinate variable operators $x = \sum_{j=1}^n \mathbf{e}_j x_j$ by its discrete counterparts

$$D_h^\pm = \Psi_\pm^\pm D (\Psi_\pm^\pm)^{-1}, \quad \text{and} \quad x^\pm = \Psi_\pm^\pm x (\Psi_\pm^\pm)^{-1} = \sum_{j=1}^n \mathbf{e}_j x_j \tau_{\mp h \mathbf{v}_j}, \quad (59)$$

respectively (see [19]).

A further consequence, we recover the Fischer Decomposition obtained in [25]. If Π_k denote the space of all Clifford-Valued polynomials homogeneous of degree k , and Π corresponds to the graded algebra

$$\Pi = \bigoplus_{k=0}^{\infty} \Pi_k,$$

the set

$$\Pi^\mp = \Psi_{\underline{x}}^\pm \Pi_k = \{f \in \Pi : E_h^\mp f = kf\}.$$

corresponds to the spaces of Clifford-valued polynomials generated by the polynomials $(\underline{x})_{\mp}^{(\alpha)}$ of degree $|\alpha| = k$, where $E_h^\mp = \sum_{j=1}^n x_j \partial_h^{\mp j}$ stands the difference Euler operators introduced in [25].

Then the Fischer Decomposition for the Difference Dirac Operators $D_h^\pm$ corresponds to

$$\Pi_k^\mp = \bigoplus_{j=0}^k (x^\pm)^j \mathcal{M}_{k-j}^\mp, \quad \text{where } \mathcal{M}_{k-j}^\mp := \Pi_k^\mp \cap \ker D_h^\pm. \quad (60)$$

We have to stress that the Fischer Decomposition obtained above does not allow to obtain a refinement of the discrete spherical harmonics (i.e., the null solutions of the star Laplacian/D'Alembertian (34)) in terms of null solutions of $D_h^\pm$. Now we will turn our attention for the discrete case described in Example 3.2.

Example 6.2 (Refinement of discrete spherical harmonics) *We shall now consider the central difference Dirac operator acting in lattices with mesh-size $h/2$, i.e.*

$$D_{h/2} := \frac{1}{2} (D_{h/2}^+ + D_{h/2}^-).$$

In this way, taking the central finite differences and the resulting inverse for the Pincherle derivative described in Example 2.2, we build the Sheffer map

$$\Psi_{\underline{x}} : \underline{x}^\alpha \mapsto \prod_{j=1}^n (x'_j)^{\alpha_j} \mathbf{1}, \quad (61)$$

viz the symmetrized formula (20) for the operators x'_j and hence, we obtain like in Example 6.1 the Almansí-type decomposition as well as the Fischer decomposition for the central difference Dirac operator $D_{h/2}$. The main advantage of the described approach we split the star Laplacian operator as the square of $D_{h/2}$.

Now we can construct the discrete counterpart of the spherical harmonics into the null solutions of $D_{h/2}$. This is essentially obtained in the following way.

Namely, if f is harmonic on the unit ball $\mathbb{B} := \{x : -x^2 \leq 1\}$, there exists a unique decomposition (see [19], page 1545)

$$f(x) = f_1(x) + x f_0(x) \quad (62)$$

where f_1 and f_0 are monogenic in $\mathbb{B}$. Moreover, f_1 and f_0 are explicitly given by

$$f_1(x) = (\mathbf{id} + x I_{n/2} D) f(x), \quad f_0(x) = -I_{n/2} D f(x)$$

Since the Sheffer map is an algebra isomorphism, we obtain the decomposition of discrete harmonic functions on the discrete ball

$$\Psi_{\underline{x}} \mathbb{B} := \left\{ x_{h/2} = \Psi_{\underline{x}} x : -(x_{h/2})^2 \mathbf{1} \leq 1 \right\},$$

by applying the Sheffer map $\Psi_{\underline{x}}$ on both sides of (62). Indeed we have

$$\Psi_{\underline{x}} f(x) = \Psi_{\underline{x}} f_1(x) + x_{h/2} \Psi_{\underline{x}} f_0(x). \quad (63)$$

Moreover if we restrict the above decomposition to the discrete sphere $\partial(\Psi_{\underline{x}} \mathbb{B})$, $\Psi_{\underline{x}} f(x) \in \Psi_{\underline{x}} \Pi_k$ is the discrete counterpart of the k -spherical harmonic polynomial $f(x)$ while $\Psi_{\underline{x}} f_1(x) \in \Psi_{\underline{x}} \Pi_k$ is the discrete counterpart of the k -spherical monogenic polynomial $f_1(\underline{x})$.

From the definition, we can also see that contrary to the operators $D_h^\pm$, the central difference Dirac operator $D_{h/2}$ is not a ‘local operator’ since the shift operators $\tau_{\pm \frac{h}{2} \mathbf{v}_j}$ when acting on some lattice functions, they not only concern with the nearest neighbor points but also all the points contained in the direction $h\mathbf{v}_j$. So we can call the combination $D_{h/2}$ a ‘quasi-local’ operator. The quasi-local is the price for getting the above mentioned nice description of discrete spherical monogenics.

The next important example concerns the Gauged version of Harmonic Oscillator written in terms of the operators described in the above example.

Example 6.3 (The gauged version of the Quantum Harmonic Oscillator) *Now we consider the Gauged version of the Quantum Harmonic Oscillator.*

$$\mathcal{H}_h = \Psi_{\underline{x}} \mathcal{H} \Psi_{\underline{x}}^{-1} = \frac{1}{2} (D_{h/2})^2 - \frac{1}{2} (x_{h/2})^2.$$

This local operator is defined on the uniform lattice with mesh-width h and corresponds to a splitting in terms of ‘quasi-local’ operators.

In order to describe the eigenfunctions of the gauged Hamiltonian, we turn now our attention to the continuous harmonic oscillator described in subsection 4.1.

In the continuum, a straightforward computation shows that

$$V_\alpha(\underline{x}) = \exp(-|\underline{x}|^2) H_\alpha(\underline{x}),$$

are eigenfunctions of $\mathcal{H}$ with corresponding eigenvalue $\lambda = |\alpha| + \frac{n}{2}$, where $H_\alpha(\underline{x})$ stands the classical multivariate Hermite polynomials.

For these eigenfunctions, we also obtain following raising and lowering properties for the set of operators $\mathbf{a}_j^\pm$:

$$\mathbf{a}_j^+ V_\alpha(\underline{x}) = V_{\alpha+\mathbf{v}_j}(\underline{x}), \quad \mathbf{a}_j^- V_\alpha(\underline{x}) = \alpha_j V_{\alpha-\mathbf{v}_j}(\underline{x}). \quad (64)$$

Moreover, the exponential generating function for $V_\alpha(\underline{x})$ is given by

$$V(\underline{x}, \underline{t}) = \sum_{\alpha} \frac{V_\alpha(\underline{x})}{\alpha!} \underline{t}^\alpha = \exp \left(-\frac{1}{2} (|\underline{x}|^2 + |\underline{t}|^2) + \sqrt{2} \underline{x} \cdot \underline{t} \right).$$

The corresponding eigenfunctions and exponential generating function for the gauged Hamiltonian $\mathcal{H}_h$ are then obtained viz the Sheffer map (61). These are closely related with the Kravchuk polynomials of discrete variable introduced in [31].

Furthermore for the spaces of Clifford valued polynomials generated by $\Psi_{\underline{x}} V_\alpha(\underline{x})$ we obtain the following version of the Fischer Decomposition

$$\bigoplus_{j=0}^k (x_{h/2})^j \Psi_{\underline{x}} H_{k-j}^+(\Omega), \quad (65)$$

where $\Psi_{\underline{x}} H_{k-j}^+(\Omega) := \{f \in \Pi : \mathcal{H}_h f = (k - j + \frac{n}{2}) f\} \cap \ker (D_{h/2})_-$.

In the last example, the Fischer decomposition results as a direct consequence of the Howe dual pair technique (see [28]).

Moreover, the following theorem appears as a generalization of (65), allow us to split the eigenvectors of the Gauge Hamiltonian $\mathcal{H}_h$ in terms of null solutions for the operator $(D_{h/2})_-$.

Theorem 6.1 (Almansi-type decomposition for the Gauge Harmonic Oscillator) *Let Ω be a star-like domain in $\mathbb{R}^n$ with center 0. If f satisfies*

$$(D_{h/2})_- f(x) = 0, \quad \text{and} \quad \mathcal{H}_h f = (k + \frac{n}{2})f(x) \quad (66)$$

in $\Omega_{h/2} := \Omega \cap (h/2)\mathbb{Z}^n$, then there exist unique functions $f_1, \dots, f_k \in \ker(D_{h/2})_-$, such that

$$f(x) = f_1(x) + (D_{h/2})_+ f_2(x) + \dots + (D_{h/2})_+^{k-1} f_k(x). \quad (67)$$

Moreover the discrete monogenic functions $f_1, \dots, f_k$ are given by the the following formulas:

$$\begin{aligned} f_k &= Q'_{k-1}(D_{h/2})_-^{k-1} f(x) \\ f_{k-1} &= Q'_{k-2}(D_{h/2})_-^{k-2} (\text{id} - (D_{h/2})_+^{k-1} Q'_{k-1}(D_{h/2})_-^{k-1}) f(x) \\ &\vdots \\ f_2 &= Q'_1(D_{h/2})_- (\text{id} - (D_{h/2})_+^2 Q'_2(D_{h/2})_-^2) \dots (\text{id} - (D_{h/2})_+^{k-1} Q'_{k-1}(D_{h/2})_-^{k-1}) f(x) \\ f_1 &= (\text{id} - (D_{h/2})_+ Q'_1(D_{h/2})_-) (\text{id} - (D_{h/2})_+^2 Q'_2(D_{h/2})_-^2) \dots (\text{id} - (D_{h/2})_+^{k-1} Q'_{k-1}(D_{h/2})_-^{k-1}) f(x). \end{aligned} \quad (68)$$

Conversely, the sum in (67), the functions $f_1, \dots, f_k$ satisfy the equations (66) in $\Omega_{h/2}$.

With the above results show the powerful Umbral bridge unifies continuous and discrete.

7 Concluding Remarks:

Umbral Calculus is an analysis of correspondence between continuity and discreteness in the commutative field related to polynomials. In the article, we bring Umbral Calculus into the non-commutative field through the Umbral Clifford analysis and we further consider functions far beyond polynomials. We study the discrete version in the Umbral Clifford analysis by keeping the symmetry of phenomena with the price of non-commutative. The Almansi decomposition is established in Umbral Clifford analysis, which concerns the theory of the powers of Laplacian. In particular, it gives the Fisher decomposition of polynomials, which is a start pointing for the spherical harmonics and special function theory as a whole. One can further study the theory of Fourier transformations and wavelets. Recently the theory of power operators of Laplacian has found their applications in the conformal Geometry in pure mathematics (see [33]).

As a framework of the unification of continuity and discreteness, the strong power of this framework lies at its constructive isomorphism. This makes it possible to construct the associated equations and solutions in the discrete setting starting from the equations and their solutions in the continuous setting in Physics. One may further bring this technique to the field of Geometry by studying the fruitful equations in the continuous setting through the associated discrete version.

References

- [1] Di Bucchanico A., Loeb, D.E. and Rota, G.C. *Umbral calculus in Hilbert space* In: B. Sagan and R.P. Stanley (eds.), *Mathematical Essays in Honor of Gian-Carlo Rota*, pp. 213-238, Birkhuser, Boston, 1998.
- [2] Roman - *The Umbral Calculus*, Academic Press, San Diego (1984)
- [3] Roman, S. and Rota, G. C. 'The umbral calculus', *Adv. Math.* 27(1978), 95-188.

- [4] Brackx, F., Delanghe and Sommen F., ‘Clifford Analysis’, Pitman Publishers, Boston-London Melbourne, 1982.
- [5] Delanghe, R., Sommen, F. and Soucek, V., Clifford algebra and spinor-valued functions. Amsterdam: Kluwer Acad. Publ., 1992.
- [6] Gilbert J. and Murray M. ‘Clifford Algebra and Dirac Operators in Harmonic Analysis,’ Cambridge University Press, Cambridge, 1991.
- [7] Gürlebeck, K., Sprössig, W., Quaternionic analysis and elliptic boundary value problems, Berlin: Akademie - Verlag, 1989.
- [8] Howe, R., Tan, E., Nonabelian harmonic analysis, Universitext. Springer-Verlag, New York, 1992. Applications of $SL(2, \mathbb{R})$ s
- [9] Lurie, S. A., Vasiliev, V. V., The Biharmonic Problem in the Theory of Elasticity, Amsterdam: Gordon and Breach Pub. 1995, 1–265.
- [10] Andersson, L. E., Elfving, T., Golub G. H., Solution of biharmonic equations with application to radar imaging, J. Comput. Appl. Math. 1998, 94(2): 153–180.
- [11] Armitage, D., Gardiner, St., Haussmann, W., Characterization of best harmonic and subharmonic L_1 -approximation, J. Reine Angew. Math. 1996, 478: 1–15.
- [12] Kounchev, O. I., Multivariate Polysplines: Applications to Numerical and Wavelet Analysis, San Diego: Academic Press 2001, 1–498.
- [13] Almansi, E., ‘Sulle integrazione dell’ equazione differenziale $\Delta^{2m}u = 0$, Ann. Mat. Pura Appl. Suppl. 3, 2., (1898).
- [14] Aronszajn, N., Creese, T. M. and Lipkin, L. J., Polyharmonic functions, Oxford Mathematical Monographs, Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1983.
- [15] Cohen J. M., Colonna F., Gowrisankaran K., and Singman D., Polyharmonic functions on trees, Amer. J. Math 124(2002), 999–1043.
- [16] Ren, G, ‘Almansi decomposition for Dunkl operators’, Science in China 48A Supp.(2005), 333–342.
- [17] Ren, G and Kähler U., ‘Almansi decomposition for polyharmonic, polyheat, and polywave functions’, Studia Math. 172(2006), 91–100.
- [18] Ren, G. and Malonek, H. ‘Decomposing kernels of iterated Operators-a unified approach’ , Math. Meth. Appl. Sci. 30(2007), 1037–1047.
- [19] Malonek, H. and Ren, G, ‘Almansi type theorems in Clifford analysis’, Math. Meth. Appl. Sci. 25(2002), 1541–1552.
- [20] Joni S. and Rota G-C - *Coalgebras and algebras in combinatorics*, Stud. Appl. Math., 61, (1979), pp. 93–139
- [21] Di Bucchianico A. and Loeb, D.E. *Operator expansion in the derivative and multiplication by x*, Integral Transf. Spec. Func. 4 (1996), 49–68.
- [22] Rota G-C, Kahane D and Odlyzko A - *Finite operator calculus*, J. Math. Anal. Appl 42 (1973)
- [23] Sommen F. *An Algebra of Abstract vector variables*, Portugaliae Math. 54, 3 (1997), 287–310
- [24] Vaz J. - *Clifford-like Calculus over Lattice*, Adv. Appl. Clifford Alg. 7, No. 1 (1997) pp. 37–70
- [25] N. Faustino, U. Kähler, *Fischer Decomposition for Difference Dirac Operators*, Adv. Appl. Cliff. Alg., 17, no. 1 (2007), 37–58.

- [26] Faustino N., *Rediscovering Clifford Analysis: On the Interplay between Umbral Calculus and Quantum Mechanics*, in preparation
- [27] Frappat L., Sciarrino A., Sorba P.- *Dictionary of Lie algebras and super algebras*, Academic Press, New York, 2000
- [28] Howe R.- *Dual pairs in physics: harmonic oscillators, photons, electrons, and singletons*, in M. Flato et al. (Eds), *Applications of groups theory in physics and mathematical physics*, AMS (1985) pp.179-208
- [29] Dimakis A., Mueller-Hoissen F., Striker T.- *Umbral calculus, discretization, and quantum mechanics on a lattice*, J. Phys. A 29 (1996) pp. 6861-6876
- [30] Delanghe, R., Brackx, F., ‘Hypercomplex function theory and Hilbert modules with reproducing kernel’, *Proc. London Math. Soc.* (3) **37**, No. 3, 545–576, (1978).
- [31] M. Lorente, *Continuous vs. discrete models for the quantum harmonic oscillator and the hydrogen atom*, Phys. Letters A, **285**, no. 1 (2001), 119-126.
- [32] Mullin R. and Rota G-C - *On the foundations of combinatorial theory III. Theory of binomial enumeration*, In Harris, editor, *Graph theory and its applications*, pp. 167-213, Academic Press, 1970
- [33] Holland J. E. and Sparling G. A. J., *Conformally invariant power of the ambient Dirac operators*, Arxiv 2001.